



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

# Ectopic Pregnancy and Epithelial to Mesenchymal Transition: is there a Link?

### Citation for published version:

Flanagan, H, Lin, C-J, Campbell, L, Horner, P, Horne, AW & Spears, N 2020, 'Ectopic Pregnancy and Epithelial to Mesenchymal Transition: is there a Link?', *Reproduction*. <https://doi.org/10.1530/REP-20-0542>

### Digital Object Identifier (DOI):

[10.1530/REP-20-0542](https://doi.org/10.1530/REP-20-0542)

### Link:

[Link to publication record in Edinburgh Research Explorer](#)

### Document Version:

Peer reviewed version

### Published In:

Reproduction

### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



**Ectopic Pregnancy and Epithelial to Mesenchymal Transition: is there a Link?**

**Authors:** Heather Flanagan<sup>1,2</sup>, Chih-Jen Lin<sup>2</sup>, Lisa L Campbell<sup>2</sup>, Paddy Horner<sup>3,4</sup>, Andrew W Horne<sup>2\*</sup>, Norah Spears<sup>1\*</sup>

**Affiliations:**

<sup>1</sup>Biomedical Sciences, University of Edinburgh, Edinburgh, Scotland

<sup>2</sup>MRC Centre for Reproductive Health, University of Edinburgh, Edinburgh, Scotland

<sup>3</sup>Population Health Sciences, University of Bristol, Bristol, UK

<sup>4</sup>NIHR Health Protection Research Unit in Behavioural Science and Evaluation at University of Bristol

\* both authors contributed equally to the work

**Corresponding authors:**

Norah Spears - [norah.spears@ed.ac.uk](mailto:norah.spears@ed.ac.uk); Andrew Horne - [andrew.horne@ed.ac.uk](mailto:andrew.horne@ed.ac.uk); Heather Flanagan - [H.C.Flanagan@sms.ed.ac.uk](mailto:H.C.Flanagan@sms.ed.ac.uk)

## **Abstract**

Ectopic pregnancy (EP) is defined as the implantation of an embryo outside of the uterus and is a leading cause of first trimester maternal mortality and morbidity. This article discusses a possible role for epithelial to mesenchymal transition in the pathogenesis of EP, given the notable similarity of protein expression between the two processes.

## **Introduction**

Ectopic pregnancy (EP) is defined as the implantation of an embryo in locations outside of the uterine cavity (Sivalingam *et al.*, 2011). EP occurs in 1-2% of all pregnancies and is a leading cause of first-trimester death worldwide. The most common implantation site for an EP is the Fallopian tube, commonly referred to as a tubal ectopic pregnancy (tEP) (Sivalingam *et al.*, 2011). Multiple risk factors predispose to tEP, such as smoking, but a unifying mechanism has not yet been identified. Current research on tEP suggests that increased embryo-receptivity is secondary to an abnormal tubal microenvironment, but there are still gaps in the literature and more research is necessary to understand the processes involved in the development of a tEP.

Epithelial to mesenchymal transition (EMT) is a biological process whereby epithelial cells undergo loss of polarity and cell-cell adhesion, then adopt mesenchymal characteristics such as migration, invasion and resistance to apoptosis (Bilyk *et al.*, 2017). EMT is associated with normal functions such as tissue regeneration and embryogenesis but the full role of EMT in various organs are not yet fully understood. Endometrial EMT is required for embryo implantation in normal intrauterine pregnancy (Bilyk *et al.*, 2017). EMT has also been associated with disease states in the reproductive tract such as ovarian cancer and adenomyosis (Bilyk *et al.*, 2017).

Here, we put forward the hypothesis that tEP occurs as a result of EMT in the epithelial cells lining the Fallopian tube, with these changes enabling ectopic embryo implantation.

## **Link between tEP and EMT**

There are many similarities in protein expression between EMT and tEP including:  $\beta$ -integrins, Wnt, Mucin-1 (MUC-1), E-cadherin, signal transducer and activator of transcription

(STAT3), and matrix metalloproteases (MMPs) (Shaw *et al.*, 2010; Bilyk *et al.*, 2017). Initial induction of EMT is under control of several factors. The upregulation of  $\beta$ 1-integrin through the TGF- $\beta$  pathway is essential for the onset of EMT (Yeh *et al.*, 2010). Wnt signalling is also described as an initial inducer of EMT, releasing  $\beta$ -catenin from E-cadherin which acts as a downstream factor increasing the transcription of EMT related proteins (Bilyk *et al.*, 2017). Cell adhesion proteins play a significant role in EMT, with a loss of cell-adhesion caused by decreased expression of MUC-1 and E-cadherin (Guaita *et al.*, 2002; Bilyk *et al.*, 2017). As EMT progresses, the basement membrane degrades, while proliferation and invasion are promoted. These events are facilitated by the upregulation of matrix metalloproteases (MMPs) and STAT3, both of which promote proliferation and invasion during EMT (Bilyk *et al.*, 2017).

Similar to the process of EMT,  $\beta$ 1-integrin is upregulated in endometrial luminal epithelium during implantation in intrauterine pregnancy, and has also been noted to be increased in the cytoplasm of Fallopian tube epithelial cells in tEP (Shaw *et al.*, 2010; Jiang *et al.*, 2019). Wnt plays an essential role in embryo attachment in endometrial luminal epithelial cells and Wnt upregulation has been shown to increase trophoblast attachment in Fallopian tubal epithelial cells in-vitro (Kodithuwakku *et al.*, 2012). Proteins affecting cell adhesion have a major role in tEP (Shaw *et al.*, 2010). In the endometrial luminal epithelium, the 'anti-adhesive' MUC1 is downregulated during implantation, enabling embryo adhesion: likewise, in tEP, there is a marked reduction in MUC-1 expression in epithelial cells of the Fallopian tube, which again could facilitate attachment of embryos due to its 'anti-adhesive' action (Shaw *et al.*, 2010). There is also a loss of E-cadherin in tEPs (Shaw *et al.*, 2010; Jiang *et al.*, 2019). MMPs play an important role in normal implantation in the extracellular matrix of the uterine luminal epithelium and tEP has been shown to have a similar MMPs expression milieu in the epithelial and smooth muscle cells of the Fallopian tube at the site of implantation (Qiu *et al.*, 2011). Furthermore, the STAT3 pathway, expressed in the uterine luminal epithelial cells during normal implantation, can be activated by leukaemia inhibitory factor, LIF (Krishnan *et al.*, 2013). LIF exposure increases trophoblast spheroid adhesion in

Fallopian tube cells in-vitro (Krishnan *et al.*, 2013): LIF also promotes EMT by activating the STAT3 pathway (Yue *et al.*, 2015). Not all EMT protein expression changes are mirrored by alterations during tEP: the SLIT/ROBO genes, described as tumour suppressor genes, are involved in the EMT process (Duncan *et al.*, 2010), and endometrial SLIT/ROBO expression has been noted to change in a temporal and spatial manner across the menstrual cycle, but examination of Fallopian tube epithelial cells, has found no difference in expression of SLIT/ROBO between samples with and without tEP (Duncan *et al.*, 2010). Other markers involved in the EMT process, such as Slug, Snail, and Twist, are expressed in the endometrium but have not yet been analysed in the Fallopian tube epithelium or tEP. Although, the link between tEP and EMT is marked, it is non-the-less possible that EMT is a consequence rather than a cause of the presence of tEP, a phenomenon that would also be of interest.

#### **Fallopian tube secretory cell expansion and EMT**

The Fallopian tube epithelium is comprised of two different cell types, secretory cells and ciliated cells. Fallopian tube secretory cells are the cells of origin for ovarian carcinoma, the most common cancer in the female pelvic organs: these secretory cells have a unique biology, co-expressing both epithelial and mesenchymal markers (Yamamoto *et al.*, 2016). Secretory cell outgrowth (SCOUT) refers to secretory cell expansion of at least 30 cells of secretory epithelial type, creating a lesion of only secretory cells compared to a mosaic of ciliated and secretory cells in normal Fallopian tubes (Yamamoto *et al.*, 2016). It is widely accepted that the origin of EMT for ovarian carcinoma is derived from the Fallopian tube secretory cells, suggesting that an altered Fallopian tube cellular structure and environment could be present years before an ovarian cancer diagnosis. Furthermore, there have been reports of the risk of ovarian carcinoma being increased in women with a history of tEP, although this is more controversial, with other literature failing to find such a link (Stewart *et al.*, 2020).

Although, SCOUT's have been researched in ovarian carcinoma; the pathophysiology and implications on tubal factor fertility of SCOUT's have not yet been determined. SCOUTS have been recently demonstrated to have upregulated LEF1 (Schmoeckel *et al.*, 2017). LEF1 activates the transcription of hallmark EMT effectors including N-Cadherin, Vimentin, and Snail (Santiago *et al.*). SCOUTS can be also found throughout the Fallopian tube (see figure 1A in Schmoeckel *et al.*, 2017). Yamamoto. *et al.*, (2016) describe an altered tubal epithelial cell microenvironment associated with immortalisation in SCOUTs and suggest that these aberrations in gene expression may even occur prior to and in the absence of serous carcinoma (Yamamoto *et al.*, 2016). When secretory cells become neoplastic, they can exhibit EMT characteristics such as increased proliferation, a loss of nuclear polarity and the ability to migrate (Yamamoto *et al.*, 2016). Additionally, tEP is associated with a loss of ciliation, and so embryo implantation in tEP cases may be proceeding at sites where that cilia loss has occurred (Shaw *et al.*, 2010; Yamamoto *et al.*, 2016). Therefore, it could be hypothesised that EMT occurring at the site of SCOUTs may lead to an ectopic implantation in the Fallopian tube.

### **Smoking and tEP**

Cigarette smoking increases the risk of tEP to 1.7-3.9% of pregnancies in smokers (Sivalingam *et al.*, 2011). In line with our hypothesis, cigarette smoke is widely associated with EMT, including having been shown to induce EMT in the reproductive system. Nicotine increases EMT in human ovarian carcinomas (Jeon *et al.*, 2016). Cotinine is an active metabolite of nicotine. Exposure of a Fallopian tube epithelial cell line to cotinine causes decreased Bcl2-associated agonist of cell death (BAD) expression, leading to decreased Bcl2 expression; reduced BAD expression is also implicated in EMT (Horne *et al.*, 2014). Another cigarette smoke component, Benzo(a)Pyrene (BaP), is known to inhibit endometrial cell apoptosis leading to impaired endometrial function. It has not been investigated if this impaired function is secondary to an EMT process but given the ability of BaP to induce EMT

141 in colonic epithelium by altering the Wnt/ $\beta$ -catenin pathway: it would seem possible (Yi *et al.*,  
142 2019). To date, the effects of BaP exposure on the Fallopian tube epithelium has not yet  
143 been researched. We hypothesise that the components of cigarette smoke may induce EMT  
144 in Fallopian tube epithelium, explaining the increased risk of tEP associated with cigarette  
145 smoking.

## 147 **Conclusion**

148  
149 In summary, EMT is likely a common process in the Fallopian tube, which could explain the  
150 pathophysiology of EP, providing a unifying a mechanism behind multiple risk factors for  
151 tEP. There are similarities in protein expression between EMT and tEP, including integrins,  
152 Wnt, MUC-1, E-cadherin, STAT3 and MMPs. There is also evidence to suggest EMT occurs  
153 in the secretory cell outgrowths of the Fallopian tube causing progression to ovarian  
154 carcinoma, although the implications on fertility and reproductive consequences in relation to  
155 ectopic pregnancy are largely unknown. Finally, there is a significant link between cigarette  
156 smoke components and EMT in the female reproductive system and we hypothesise this  
157 may extend to the Fallopian tube epithelium. In conclusion, the similarities in the literature  
158 between EMT and tEP suggests that further research is required to understand if there is a  
159 clear link between tEP and EMT, and if so, whether EMT is a causative factor in tEP.

## 161 **Funding**

162  
163 This work was supported by a UK Medical Research Council Centre Grant (to AH, NS, JCL,  
164 LC; MR/N022556/1, 2018-22) and a joint Medical Research Council / Ectopic Pregnancy  
165 Trust PhD Fellowship to HF.

## 167 **Acknowledgments**



169 The authors thank Beckie Doust for graphical design.

170

171 **Authors' Roles**

172

173 HF conceived and drafted the manuscript; CJL, LLC and PH helped draft the manuscript;

174 AWH and NS conceived and helped draft the manuscript. All authors read and approved the

175 final version of the manuscript.

176

177 **Declaration of Interests**

178 The authors declare that there is no conflict of interest that could be perceived as prejudicing

179 the impartiality of the research reported.

180

181

## References

- Bilyk O, Coatham M, Jewer M and Postovit L-M (2017) Epithelial-to-Mesenchymal Transition in the Female Reproductive Tract: From Normal Functioning to Disease Pathology. *Frontiers in Oncology* **7** 145.
- Duncan WC, McDonald SE, Dickinson RE, Shaw JLV, Lourenco PC, Wheelhouse N, Lee K-F, Critchley HOD and Horne AW (2010) Expression of the repulsive SLIT/ROBO pathway in the human endometrium and Fallopian tube. *Molecular Human Reproduction* **16** 950–959.
- Guaita S, Puig I, Francí C, Garrido M, Domínguez D, Batlle E, Sancho E, Dedhar S, Herreros AG de and Baulida J (2002) Snail Induction of Epithelial to Mesenchymal Transition in Tumor Cells Is Accompanied by MUC1 Repression and ZEB1 Expression. *Journal of Biological Chemistry* **277** 39209–39216.
- Horne AW, Brown JK, Nio-Kobayashi J, Abidin HBZ, Adin ZEHA, Boswell L, Burgess S, Lee K-F and Duncan WC (2014) The Association between Smoking and Ectopic Pregnancy: Why Nicotine Is BAD for Your Fallopian Tube. *PLoS ONE* **9** e89400.
- Jeon S-Y, Go R-E, Heo J-R, Kim C-W, Hwang K-A and Choi K-C (2016) Effects of cigarette smoke extracts on the progression and metastasis of human ovarian cancer cells via regulating epithelial-mesenchymal transition. *Reproductive Toxicology* **65** 1–10.
- Jiang H, Yang X-Y and Zhu W-J (2019) Networks of E-cadherin,  $\beta$  1 integrin, and focal adhesion kinase in the pathogenesis of tubal pregnancy. *Gynecological Endocrinology* **35** 346–350.
- Kodithuwakku SP, Pang RT, Ng EH, Cheung AN, Horne AW, Ho PC, Yeung WS and Lee KF (2012) Wnt activation downregulates olfactomedin-1 in Fallopian tubal epithelial cells: a microenvironment predisposed to tubal ectopic pregnancy. *Laboratory Investigation; a Journal of Technical Methods and Pathology* **92** 256–264.
- Krishnan T, Winship A, Sonderegger S, Menkhurst E, Horne AW, Brown J, Zhang J-G, Nicola NA, Tong S and Dimitriadis E (2013) The role of leukemia inhibitory factor in tubal ectopic pregnancy. *Placenta* **34** 1014–1019.
- Qiu X, Xie Y, Chen L and Gemzell-Danielsson K (2011) Expression of matrix metalloproteinases and their inhibitors at the feto-maternal interface in unruptured ectopic tubal pregnancy. *Acta Obstetrica et Gynecologica Scandinavica* **90** 966–971.
- Santiago L, Daniels G, Wang D, Deng F-M and Lee P Wnt signaling pathway protein LEF1 in cancer, as a biomarker for prognosis and a target for treatment. 18.
- Schmoeckel E, Odai-Afotey AA, Schleißheimer M, Rottmann M, Flesken-Nikitin A, Ellenson LH, Kirchner T, Mayr D and Nikitin AY (2017) LEF1 is preferentially expressed in the tubal-peritoneal junctions and is a reliable marker of tubal intraepithelial lesions. *Modern Pathology* **30** 1241–1250.
- Shaw JLV, Dey SK, Critchley HOD and Horne AW (2010) Current knowledge of the aetiology of human tubal ectopic pregnancy. *Human Reproduction Update* **16** 432–444.
- Sivalingam VN, Duncan WC, Kirk E, Shephard LA and Horne AW (2011) Diagnosis and management of ectopic pregnancy. *Journal of Family Planning and Reproductive Health Care* **37** 231–240.

221 Stewart LM, Stewart CJR, Spilsbury K, Cohen PA and Jordan S (2020) Association between pelvic  
 222 inflammatory disease, infertility, ectopic pregnancy and the development of ovarian serous  
 223 borderline tumor, mucinous borderline tumor and low-grade serous carcinoma. *Gynecologic*  
 224 *Oncology* **156** 611–615.

225 Yamamoto Y, Ning G, Howitt BE, Mehra K, Wu L, Wang X, Hong Y, Kern F, Wei TS, Zhang T *et al.*  
 226 (2016) *In vitro* and *in vivo* correlates of physiological and neoplastic human Fallopian tube  
 227 stem cells: *In vitro* and *in vivo* correlates of physiological and neoplastic human FTSCs. *The*  
 228 *Journal of Pathology* **238** 519–530.

229 Yeh Y-C, Wei W-C, Wang Y-K, Lin S-C, Sung J-M and Tang M-J (2010) Transforming Growth Factor- $\beta$ 1  
 230 Induces Smad3-Dependent  $\beta$ 1 Integrin Gene Expression in Epithelial-to-Mesenchymal  
 231 Transition during Chronic Tubulointerstitial Fibrosis. *The American Journal of Pathology* **177**  
 232 1743–1754.

233 Yi T, Liu M, Li X, Liu X, Ding Y, He J, Xu H, Gao R, Mu X, Geng Y *et al.* (2019) Benzo(a)pyrene inhibits  
 234 endometrial cell apoptosis in early pregnant mice via the WNT5A pathway. *Journal of*  
 235 *Cellular Physiology* **234** 11119–11129.

236 Yue X, Zhao Y, Zhang C, Li J, Liu Z, Liu J and Hu W (2015) Leukemia inhibitory factor promotes EMT  
 237 through STAT3-dependent miR-21 induction. *Oncotarget* **7** 3777–3790.

238  
 239  
 240  
 241  
 242

## **Figure legend**

**Figure 1. Similarities between EMT, EP and intrauterine pregnancy.** Epithelial cells displaying polarity are held together by adhesion proteins and tight junctions. Induction of EMT effector proteins such as integrins and Wnt signalling leads to a downstream loss of cell adhesion proteins; blue and pink receptors represent MUC-1 and E-Cadherin receptors respectively: scissors represent a loss of these proteins. Due to the loss of cell adhesion and a loss of microvilli, epithelial cells resort to a mesenchymal like state. This leads to the upregulation of STAT3 and MMPs which results in basement membrane degradation and induces proliferation and invasion of mesenchymal cells which are loosely arranged in the extracellular matrix (ECM). Both EP and intrauterine pregnancies have similar protein expression patterns to those expressed in the EMT process.

